

Gulf of Mexico Harmful Algal Bloom Bulletin

Region: Texas


Monday, 25 January 2016

NOAA National Ocean Service

NOAA Satellite and Information Service

NOAA National Weather Service

Last bulletin: Tuesday, January 19, 2016



Satellite chlorophyll image with possible *K. brevis* HAB areas shown by red polygon(s), when applicable. Points represent cell concentration sampling data from January 15 to 22: red (high), orange (medium), yellow (low b), brown (low a), blue (very low b), purple (very low a), pink (present), and green (not present). Cell count data are provided by Texas Parks and Wildlife Department. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide:

http://tidesandcurrents.noaa.gov/hab/habfs_bulletin_guide.pdf

Detailed sample information can be obtained through the Texas Parks and Wildlife Department at: http://www.tpwd.state.tx.us./landwater/water/environconcerns/hab/redtide/status.phtml

Conditions Report

Karenia brevis (commonly known as Texas red tide) ranges from not present to very low concentrations along the coast of Texas. No respiratory irritation is expected Monday, January 25 through Monday, February 1.

Check http://tidesandcurrents.noaa.gov/hab/beach_conditions.html for recent, local observations.


Analysis

Sampling from the Texas A&M University's Imaging FlowCytobot, located on the Port Aransas ship channel, indicates that *Karenia brevis* ranges between 'not present' and 'very low a' concentrations (TAMU; 1/20-25). For information on area shellfish restrictions, contact the Texas Department of State Health Services.

In recent MODIS Aqua imagery (1/24, shown left) patches of elevated to very high chlorophyll (2 to >20 μ g/L) are visible along- and offshore the Texas coast from Sabine Pass to the Padre Island National Seashore region. Elevated chlorophyll is not indicative of the presence of *K. brevis* and is most likely due to the resuspension of benthic chlorophyll and sediments along the coast.

Forecast models based on predicted near-surface currents indicate a potential maximum transport of 10 km south from the Port Aransas region from January 24-28.

Yang, Davis




Wind speed and direction are averaged over 12 hours from buoy measurements. Length of line indicates speed; angle indicates direction. Red indicates that the wind direction favors upwelling near the coast. Values to the left of the dotted vertical line are measured values; values to the right are forecasts. Wind observation and forecast data provided by NOAA's National Weather Service (NWS).

Wind Analysis

Port Aransas to Baffin Bay: South to southeast winds (5-20kn, 3-10m/s) today. Northeast to northwest winds (5-25kn, 3-13m/s) Tuesday through Thursday afternoon. Southwest to south winds (5-15kn, 3-8m/s) Thursday evening through Friday.

To see previous bulletins and forecasts for other Harmful Algal Bloom Bulletin regions, visit the NOAA Harmful Algal Bloom Operational Forecast System bulletin archive: http://tidesandcurrents.noaa.gov/hab/bulletins.html



Verified and suspected HAB areas shown in red. Other areas of high chlorophyll concentration shown in yellow (see p. 1 analysis for interpretation).